

KWS INSPIRATOR features the shortest, strongest straw available, ensuring excellent standability and smooth, efficient harvestability. This combination of high performance and durability makes it a top choice for farmers looking to maximize productivity and ease of harvest.

	KWS INSPIRATOR	Average of Hybrid Rye check varieties	Average of traditional check varieties
Yield (%)	100 %	94 %	72 %
Winter Survival (%)	93	93	94
Height (cm)	82	85	98
Loging (1-9)	1.5	1.6	2.0

^{*} Source: Fall Rye Western Cooperative Registration Trials 2020-21, 2021-22, 2022-23

Main Features

Market-leading yields:

- 6% over hybrid rye checks*
- 39 % over traditional checks*

Variety Highlights

- High Yield
- Excellent ergot protection
- Market-leading standability

End Use Markets

- Food
- Feed
- Distilling
- KWS COVER+





KWS RyeGHT Seeding Practices

RyeGHT Rate and Time

The standard seeding rate is 1 unit/acre under good conditions for grain or forage use. To establish hybrid fall rye successfully, three weeks of good growing conditions are needed.

Seeding rate decision factors include moisture and heat units at seeding time, trash penetration, and seed row spacing.

The suggested seeding rate and target shoot growth are as follows:

Suggested Seeding Rates

Earlier Seeding

Units/acre	1.0-1.2	
Target Shoot Growth	4-6 inches	
Later Seeding		
Units/acre	1.0-1.4	

Please consult provincial crop insurance deadlines for insurable planting dates.

RyeGHT Preparation

Fall is busy, and fall seeded crops require proactive planning. Select well drained fields that are free of residual herbicides. Know where you plan to seed your hybrid rye before seeding your spring crop, seed that spring crop early, and spread the trash evenly during harvest to limit residue issues. Have your drill cleaned and ready after spring seeding is complete.

RyeGHT Seed Placement

Seed shallow at a ¾" depth with a travel speed of <4.5 mph into a firm seed bed to maintain consistent seed depth. Seeding deeper or faster will result in higher seedling mortality and a reduced stand. Maintain a ¾" seed depth even if the soil is dry and wait for rain. Narrow row spacing of 6–7.5" is optimal for hybrid fall rye. If seeding on 10–12" spacing, speed, depth, and rate are more critical to eliminate skips and gaps in the field.

RyeGHT Stand

Plant stand target is 16-18 plants/ft² (grain) and 16-20 plants/ft² (forage). Even emergence and achieving the target stand will provide the most consistent yields and the best resistance against ergot.

RyeGHT Pest Control

To protect seedlings against disease and insects, hybrid fall rye should be professionally treated with a registered seed treatment. A pre-plant or pre-emergent herbicide application is important to control weeds when the hybrid rye is the least competitive. If weed pressure is high in spring, an in crop herbicide application may be required. A well established hybrid rye stand will be competitive with most spring emerging weeds.

Laura Eastwood, Ph.D.

Animal Nutrition Manager 368.999.1790 laura.eastwood@kws.com



Kenny Piecharka

Eastern Canada 403.308.4900 kenny.piecharka@kws.com