

KWS SANDOR

Top in Class for Ergot

- For farmers seeking reliable yields and minimal ergot risk.



KWS SANDOR is your top choice for superior ergot protection and high grain yields. This hybrid delivers outstanding performance for food, feed, and fuel markets.

Main Features

Market leading yields:

- 27 % less ergot than hybrid checks*
- 29 % less ergot than traditional checks*

Variety Highlights

- Top-in-class ergot protection
- High grain yields
- Short straw

End Use Markets

- All Grain Markets
- Feed
- Forage
- KWS **COVER+**[™]
soil protection program

	KWS SANDOR	Average of Hybrid Rye check varieties	Average of traditional check varieties
Yield (%)	100 %	101 %	76 %
Winter Survival (%)	89	88	89
Height (cm)	85	86	100
Logging (1-9)	3.0	2.3	3.5
Test Weight (kg/hL)	73.7	74.3	74.5

* Source: Fall Rye Western Cooperative Registration Trials 2018-19, 2019-20, 2020-21*

Juel Scott

Western Prairies
825.288.8861
juel.scott@kws.com



www.kws.com/ca/en

Dan Bolton

Eastern Prairies
204.595.9154
daniel.bolton@kws.com



KWS RyeGHT Seeding Practices

RyeGHT Rate and Time

The standard seeding rate is **1 unit/acre** under good conditions for grain or forage use. To establish hybrid fall rye successfully, three weeks of good growing conditions are needed.

Seeding rate decision factors include moisture and heat units at seeding time, trash penetration, and seed row spacing.

The suggested seeding rate and target shoot growth are as follows:

Suggested Seeding Rates

Earlier Seeding	
Units/acre	1.0–1.2
Target Shoot Growth	4–6 inches
Later Seeding	
Units/acre	1.0–1.4
Target Shoot Growth	3–4 inches

* Please consult provincial crop insurance deadlines for insurable planting dates.

RyeGHT Preparation

Fall is busy, and fall seeded crops require proactive planning. Select well drained fields that are free of residual herbicides. Know where you plan to seed your hybrid rye before seeding your spring crop, seed that spring crop early, and spread the trash evenly during harvest to limit residue issues. Have your drill cleaned and ready after spring seeding is complete.

RyeGHT Seed Placement

Seed shallow at a **¾" depth** with a travel **speed of <4.5 mph** into a firm seed bed to maintain consistent seed depth. Seeding deeper or faster will result in higher seedling mortality and a reduced stand. Maintain a ¾" seed depth even if the soil is dry and wait for rain. Narrow row spacing of 6–7.5" is optimal for hybrid fall rye. If seeding on 10–12" spacing, speed, depth, and rate are more critical to eliminate skips and gaps in the field.

RyeGHT Stand

Plant stand target is **16-18 plants/ft²** (grain) and **16-20 plants/ft²** (forage). Even emergence and achieving the target stand will provide the most consistent yields and the best resistance against ergot.

RyeGHT Pest Control

To protect seedlings against disease and insects, hybrid fall rye should be professionally treated with a registered seed treatment. A pre-plant or pre-emergent herbicide application is important to control weeds when the hybrid rye is the least competitive. If weed pressure is high in spring, an in crop herbicide application may be required. A well established hybrid rye stand will be competitive with most spring emerging weeds.

Juel Scott

Western Prairies
825.288.8861
juel.scott@kws.com



www.kws.com/ca/en

Dan Bolton

Eastern Prairies
204.595.9154
daniel.bolton@kws.com